

# Wide Usage Area

Dr. Clot Savior hemostat is designed to stop bleeding in capillary bleeding, parenchymatous organ bleeding, and bleeding in reaction areas during surgical operations. This hemostat is used when other hemostatic procedures are non-applicable. It has wide usage area. It can be used in important surgical operations



**General  
Surgery**



**Digestive System  
Surgery**



**Neurosurgery  
Operations**



**Liver  
Operations**



**Orthopaedics  
Operations**



**Gynaecology  
Operations**



**Urology  
Operations**



**Stomatology  
Operations**



**Throat  
Operations**



**Nasal  
Operations**



**Traumatology  
Surgery**



**Cardiovascular  
Surgery**



**Face and Jaw  
Surgery**



**Lung  
Operations**



**Gall Bladder  
Operations**

**Biopsy  
Operations**

This haemostat is suitable for all other surgical fields including general surgery, intestine system surgery, neurosurgery (especially brain operations) with restrictions, plastic surgery, orthopaedics, gynaecology, urology, stomatology, otorhinolaryngology operations, traumatology and cardiovascular operations, vascular prosthetics operations, haemorrhoidectomy, biopsy, liver and gall bladder operations, gastric resections, thoracic and abdominal sympathectomy.

Alufer-Bel absorbable hemostat can be applied to cavities (after tumour scraping) as well as endoscopic operations and dental applications.

Dr. Clot hemostat is designed to stop bleeding in capillary bleeding, parenchymatous organ bleeding, and bleeding in reaction areas during surgical operations. This hemostat is used when other hemostatic procedures are non-applicable. It has wide usage area. It can be used in important surgical operations.